

5. Sharoky CE, Sellers MM, Keele LJ, et al. Does surgeon sex matter?: practice patterns and outcomes of female and male surgeons. *Ann Surg* 2018; **267**: 1069–76
6. Sarsons H, Akhtari M, Barron K, et al. Interpreting signals in the labor market: evidence from medical referrals. Available from: https://scholar.harvard.edu/files/sarsons/files/sarsons_jmp.pdf. [Accessed 11 December 2017]
7. Zegers M, de Bruijne MC, de Keizer B, et al. The incidence, root-causes, and outcomes of adverse events in surgical units: implication for potential prevention strategies. *Patient Saf Surg* 2011; **5**: 13. <https://doi.org/10.1186/1754-9493-5-13>
8. Gonzales G, Ehrenfeld JM. Sex is not gender and why it matters for population health. *Br J Anaesth* 2018; **120**: 1130–1
9. Coates J. *Women, men and language: a sociolinguistic account of gender differences in language*. New York, NY: Routledge; 2016
10. Garden AL, Weller JM. Speaking up: does anaesthetist gender influence teamwork and collaboration? *Br J Anaesth* 2017; **119**: 571–2
11. Amacher SA, Schumacher C, Legeret C, et al. Influence of gender on the performance of cardiopulmonary rescue teams. *Crit Care Med* 2017; **45**: 1184–91
12. Jones LK, Jennings BM, Higgins MK, de Waal FBM. Ethological observations of social behavior in the operating room. *Proc Natl Acad Sci USA* 2018; **115**: 7575–80
13. Pattni N, Bould MD, Hayter MA, et al. Gender, power and leadership: the effect of a superior's gender on respiratory therapists' ability to challenge leadership during a life-threatening emergency. *Br J Anaesth* 2017; **119**: 697–702
14. Adams TL. Gender and feminization in health care professions. *Sociol Compass* 2010; **4**: 454–65 [Online]
15. Rajacich D, Kane D, Williston C, Cameron S. If they do call you a nurse, it is always a "male nurse": experiences of men in the nursing profession. *Nurs Forum* 2013; **48**: 71–80
16. Makary MA, Sexton JB, Freischlag JA, et al. Operating room teamwork among physicians and nurses: teamwork in the eye of the beholder. *J Am Coll Surg* 2006; **202**: 746–52
17. Mills A, Neal-Smith J, Bridges D. *Absent aviators: gender issues in aviation*. New York: Routledge; 2014
18. Goldenberg MG, Jung J, Grantcharov TP. Using data to enhance performance and improve quality and safety in surgery. *JAMA Surg* 2017; **152**: 972–3

British Journal of Anaesthesia, 121 (5): 999–1001 (2018)

doi: [10.1016/j.bja.2018.07.032](https://doi.org/10.1016/j.bja.2018.07.032)

Advance Access Publication Date: 5 September 2018

© 2018 British Journal of Anaesthesia. Published by Elsevier Ltd. All rights reserved.

Ward monitoring 3.0

F. Michard^{1,*} and D. I. Sessler²

¹MiCo, Denens, Switzerland and ²Department of Outcomes Research, Anesthesiology Institute, Cleveland Clinic, Cleveland, OH, USA

*Corresponding author. E-mail: frederic.michard@bluewin.ch

"Patients who are admitted to hospital believe that they are entering a place of safety. They feel confident that, should their condition deteriorate, they are in the best place for prompt and effective treatment. Yet there is evidence to the contrary".

This alarming statement is not from the victim of medical error or a burned-out clinician; it is from the National Institute for Health & Care Excellence (NICE) in the UK.

In fact, patients often die on hospital wards where clinical deterioration can remain unnoticed for hours. In the European Surgical Outcomes Study¹ (EuSOS), which included 46 539 patients from 498 hospitals in 28 countries, most patients (73%) who died were not admitted to critical care at any stage after surgery. In a national UK audit,² among 23 554 in-hospital cardiac arrests, more than half occurred on medical or surgical wards.

Ward monitoring 1.0 or intermittent spot-checks

Most ward patients do not precipitously deteriorate. Instead, gradual decline is often detected late because vital signs are usually measured at 4–6 h intervals. A study of continuous clinician-blind oxygen saturation monitoring on surgical

wards showed that saturation was <90% for more than 1 h in a third of patients, and <90% for more than 6 h in 10% of patients.³ Concerningly, nurses who measured vital signs every 4 h missed 90% of the hypoxaemic episodes lasting at least 1 h.³ Ward hypotension is similarly common, profound, and prolonged—and even more worrisome as postoperative hypotension is associated with nearly a three-fold increase in the risk of myocardial infarction.⁴

Ward monitoring 2.0 or continuous bedside monitoring

As cardiorespiratory deterioration is frequently missed with routine vital sign monitoring, it is reasonable to anticipate that continuous monitoring will improve ward safety. There is at least some evidence to support this. Taenzer and colleagues⁵ continuously monitored heart rate and oxygen saturation in 2841 patients recovering from orthopaedic surgery, most of whom were given opioids. There were fewer rescue events and critical care transfers than in parallel control wards. Brown and colleagues⁶ used a contact-free sensor put under the mattress to continuously monitor heart rate and ventilatory frequency in 2314 medical–surgical ward patients. The

number of calls for cardiac arrest decreased significantly. A recent UK study⁷ which included 4402 ward patients reported a significant decrease in cardiac arrest and hospital mortality after implementation of a continuous Early Warning Score surveillance system that used pagers to alert nurses when patients began to deteriorate.












Ward monitoring 3.0

Medical-grade wireless and wearable sensors facilitate continuous monitoring and may improve outcomes in hospital wards. Continuous monitoring of pulse rate and arterial oxygen saturation is available from pulse oximeters, some of which are now wireless. Heart rate and ECG monitoring can be obtained from various chest adhesive patches in which two electrodes are embedded a few centimetres apart (Table 1). Some detect and diagnose cardiac arrhythmias better than classical Holter systems.⁸ Ventilatory frequency is the highest-ranked variable in mathematical models predicting clinical deterioration on the wards,⁹ and is classically measured from capnographic or acoustic signals. However, ventilatory frequency can also be derived from wireless pulse oximeters and chest patch systems that sense respiratory variation in R-wave amplitude and in RR or pulse intervals (respiratory sinus arrhythmia). Chest patches containing an accelerometer, or a piezoelectric sensor can also estimate ventilatory frequency from chest movement. Several adhesive patches contain a thermistor and measure skin temperature; some are able to accurately estimate central body temperature from the axilla.¹⁰ The finger cuff pulse-decomposition method enables continuous monitoring of blood pressure wirelessly, whereas pulse-wave transit time techniques are useful to

track changes in blood pressure (i.e. to detect hypotensive events) and trigger oscillometric brachial cuff measurements.¹¹ Necklaces equipped with bioimpedance sensors have been cleared to detect changes in thoracic fluid content and may have value to diagnose pulmonary congestion before patients develop symptoms. Untethered monitors also have potential to detect patient location, position, and activity. For example, a supine unmoving patient is expected in bed, but would be a sign of collapse in a bathroom.

Wearables may well make continuous physiologic monitoring practical for ward patients (Table 1). But continuous monitoring will also generate huge amounts of data. Furthermore, wearables seem likely to be less reliable than conventional monitors and nurses who can identify and correct obviously flawed values. Simple threshold alarms for individual data streams will almost surely generate an excessive number of false alarms, but integrated Early Warning Scores could help decrease alarm fatigue.⁷ Importantly, future monitoring systems might identify trends and patterns that indicate a failing patient before any particular value exceeds conventional risk thresholds. Decreasing ventilatory frequency and arterial oxygen saturation, for example, might suggest opioid-induced respiratory depression. Increasing ventilatory frequency, heart rate, and temperature with decreasing oxygen saturation and blood pressure may suggest pulmonary sepsis—and might do so before any individual value exceeds a conventional alarm threshold. Data integration should also be able to identify artifact in one data stream by considering normal values from other sensors. Simultaneous development of computing power and predictive analytics will surely help tame the continuous flow of physiologic

Table 1 Examples of medical-grade adhesive patches for physiologic monitoring. BP, blood pressure; HR, heart rate; PR, pulse rate; PWT, pulse wave transit time; SpO₂, peripheral capillary oxygen saturation; T, temperature

	Company website	Sensor(s)	PR/HR	ECG	Ventilatory frequency	SpO ₂	BP	T
	Intelesens.com	Chest patch (x2)	+	+	+			Skin
	Irrhythmtech.com	Chest patch	+	+				
	Isansys.com	Chest patch	+	+	+			
	Medicompinc.com	Chest patch	+	+				
	Medtronic.com	Chest patch	+	+				
	Pmd-solutions.com	Chest patch			+			
	Preventicesolutions.com	Chest patch	+	+	+			
	Railing.com	Axillary patch						Axilla
	Sensium.co.uk	Chest patch, axillary thermistor	+		+			Axilla
	Smartcardia.com	Chest patch	+	+		+	PWT	Skin
	Vitalconnect.com	Chest patch	+	+	+			Skin

information and provide clinicians with more meaningful and actionable information, less artifact, and fewer false alarms. Machine learning algorithms, for example, have been used with success to distinguish real events from artifact in online multisignal vital sign monitoring data streams.¹² They may also better predict clinical deterioration than current Early Warning Score systems.⁹

Conclusions

Avoidable deaths on hospital wards remain all too common. Many should be prevented by continuous vital sign monitoring. Wireless and wearable sensors can help as patients poorly tolerate tethered monitors.¹³ Although many technical solutions already exist to monitor vital signs wirelessly, validation studies remain scarce. Major trials are needed to determine whether wireless and wearable sensors accurately monitor vital signs, avoid excessive false alarms, detect clinical deterioration sufficiently early to allow effective intervention, and reduce serious adverse outcomes.

Authors' contributions

Writing/revising editorial: both authors.

Declaration of interest

Frederic Michard is the founder and managing director of MiCo, a Swiss consulting firm. MiCo does not sell any medical product and Frederic Michard does not own shares from any medtech company. Sotera, Inc provides equipment and funding for the Department of Outcomes Research led by Daniel I Sessler.

References

1. Pearse RM, Moreno RP, Bauer P, et al. Mortality after surgery in Europe: a 7 day cohort study. *Lancet* 2012; **380**: 1059–65
2. Nolan JP, Soar J, Smith GB, et al. Incidence and outcome of in-hospital cardiac arrest in the United Kingdom national cardiac arrest audit. *Resuscitation* 2014; **85**: 987–92
3. Sun Z, Sessler DI, Dalton JE, et al. Postoperative hypoxemia is common and persistent: a prospective blinded observational study. *Anesth Analg* 2015; **121**: 709–15
4. Sessler DI, Meyhoff CS, Zimmerman NM, et al. Period-dependent associations between hypotension during and for 4 days after noncardiac surgery and a composite of myocardial infarction and death: a substudy of the POISE-2 trial. *Anesthesiology* 2018; **128**: 317–27
5. Taenzer AH, Pyke JB, McGrath SP, et al. Impact of pulse oximetry surveillance on rescue events and intensive care unit transfers: a before-and-after concurrence study. *Anesthesiology* 2010; **112**: 282–7
6. Brown H, Terrence J, Vasquez P, et al. Continuous monitoring in an inpatient medical–surgical unit: a controlled clinical trial. *Am J Med* 2014; **127**: 226–32
7. Subbe CP, Duller B, Bellomo R. Effect of an automated notification system for deteriorating ward patients on clinical outcomes. *Crit Care* 2017; **21**: 52
8. Barrett PM, Komatireddy R, Haaser S, et al. Comparison of 24-hour Holter monitoring with 14-day novel adhesive patch electrocardiographic monitoring. *Am J Med* 2014; **127**: 95.e11–7
9. Churpek MM, Yuen TC, Winslow C, et al. Multicenter comparison of machine learning methods and conventional regression for predicting clinical deterioration on the wards. *Crit Care Med* 2016; **44**: 368–74
10. Pei L, Huang Y, Mao G, Sessler DI. Axillary temperature, as recorded by the iThermonitor WT701, well represents core temperature in adults having noncardiac surgery. *Anesth Analg* 2018; **126**: 833–8
11. Michard F, Sessler DI, Saugel B. Non-invasive arterial pressure monitoring revisited. *Intensive Care Med* 2018. <https://doi.org/10.1007/s00134-018-5108-x>. Access Published March 7
12. Chen L, Dubrawski A, Wang D, et al. Using supervised machine learning to classify real alerts and artifact in online multisignal vital sign monitoring data. *Crit Care Med* 2016; **44**: e456–63
13. Michard F, Gan TJ, Kehlet H. Digital innovations and emerging technologies for enhanced recovery programmes. *Br J Anaesth* 2017; **119**: 31–9

British Journal of Anaesthesia, 121 (5): 1001–1004 (2018)

doi: [10.1016/j.bja.2018.08.007](https://doi.org/10.1016/j.bja.2018.08.007)

Advance Access Publication Date: 21 September 2018

© 2018 British Journal of Anaesthesia. Published by Elsevier Ltd. All rights reserved.

Bedside assessment of lung aeration and stretch

A. Fernandez-Bustamante^{1,2} and M. F. Vidal Melo^{3,*}

¹Department of Anesthesiology, University of Colorado School of Medicine, Aurora, CO, USA, ²Webb Waring Center, University of Colorado School of Medicine, Aurora, CO, USA and ³Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

*Corresponding author. E-mail: VidalMelo.Marcos@mgh.harvard.edu

Knowledge on lung physiology and lung injury mechanisms related to positive pressure ventilation has evolved from

animal models advancing the concepts of volutrauma, barotrauma, atelectrauma, and biotrauma^{1,2} to applications in