

JAMA Surgery | Original Investigation

Effect of a Modified Hospital Elder Life Program on Delirium and Length of Hospital Stay in Patients Undergoing Abdominal Surgery

A Cluster Randomized Clinical Trial

Cheryl Chia-Hui Chen, RN, DNSc; Hsiu-Ching Li, RN, MSN; Jin-Tung Liang, MD, PhD; I-Rue Lai, MD, PhD; Jerry Dwi Trijoyo Purnomo, MS; Yi-Ting Yang, RN, MSN; Been-Ren Lin, MD, PhD; John Huang, MD; Ching-Yao Yang, MD, PhD; Yu-Wen Tien, MD, PhD; Chiung-Nien Chen, MD, PhD; Ming-Tsan Lin, MD, PhD; Guan-Hua Huang, PhD; Sharon K. Inouye, MD, MPH

IMPORTANCE Older patients undergoing abdominal surgery commonly experience preventable delirium, which extends their hospital length of stay (LOS).

OBJECTIVE To examine whether a modified Hospital Elder Life Program (mHELP) reduces incident delirium and LOS in older patients undergoing abdominal surgery.

DESIGN, SETTING, AND PARTICIPANTS This cluster randomized clinical trial of 577 eligible patients enrolled 377 older patients (≥ 65 years of age) undergoing gastrectomy, pancreaticoduodenectomy, and colectomy at a 2000-bed urban medical center in Taipei, Taiwan, from August 1, 2009, through October 31, 2012. Consecutive older patients scheduled for elective abdominal surgery with expected LOS longer than 6 days were enrolled, with a recruitment rate of 65.3%. Participants were cluster randomized by room to receive the mHELP or usual care.

INTERVENTIONS The intervention (implemented by an mHELP nurse) consisted of 3 protocols administered daily: orienting communication, oral and nutritional assistance, and early mobilization. Intervention group participants received all 3 mHELP protocols postoperatively, in addition to usual care, as soon as they arrived in the inpatient ward and until hospital discharge. Adherence to protocols was tracked daily.

MAIN OUTCOMES AND MEASURES Presence of delirium was assessed daily by 2 trained nurses who were masked to intervention status by using the Confusion Assessment Method. Data on LOS were abstracted from the medical record.

RESULTS Of 577 eligible patients, 377 (65.3%) were enrolled and randomly assigned to the mHELP ($n = 197$; mean [SD] age, 74.3 [5.8] years; 111 [56.4%] male) or control ($n = 180$; mean [SD] age, 74.8 [6.0] years; 103 [57.2%] male) group. Postoperative delirium occurred in 13 of 196 (6.6%) mHELP participants vs 27 of 179 (15.1%) control individuals, representing a relative risk of 0.44 in the mHELP group (95% CI, 0.23–0.83; $P = .008$). Intervention group participants received the mHELP for a median of 7 days (interquartile range, 6–10 days) and had a shorter median LOS (12.0 days) than control participants (14.0 days) ($P = .04$).

CONCLUSIONS AND RELEVANCE For older patients undergoing abdominal surgery who received the mHELP, the odds of delirium were reduced by 56% and LOS was reduced by 2 days. Our findings support using the mHELP to advance postoperative care for older patients undergoing major abdominal surgery.

TRIAL REGISTRATION clinicaltrials.gov Identifier: [NCT01045330](https://clinicaltrials.gov/ct2/show/study/NCT01045330).

JAMA Surg. 2017;152(9):827-834. doi:10.1001/jamasurg.2017.1083
Published online May 24, 2017.

← Invited Commentary
page 834

+ Supplemental content

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: Guan-Hua Huang, PhD, Institute of Statistics, National Chiao Tung University, 1001 Ta Hsueh Rd, Hsinchu 30010, Taiwan, Republic of China (ghuang@stat.nctu.edu.tw).

Patients undergoing abdominal surgery often develop delirium, which greatly influences their postoperative course of clinical recovery and length of hospital stay (LOS).¹⁻³ Delirium affects 13% to 50% of patients undergoing noncardiac surgery,⁴ and the health care costs attributable to delirium are more than \$164 billion per year in the United States.^{5,6} Older surgical patients (≥65 years of age) have a particularly high risk for developing delirium, with detrimental effects on their recovery.⁶ Delirium has been associated with alterations in cholinergic activity, inflammatory processes induced by neural signaling, and excessive depth of anesthesia and sedation.^{7,8} Delirium may also be precipitated by factors such as infection, malnutrition, electrolyte and fluid imbalances, anemia, and social isolation.^{4,9,10} Nevertheless, 30% to 40% of cases of delirium are preventable¹¹; thus, implementing effective interventions to prevent incident delirium and reduce LOS is a clinical priority.

We hypothesized that delirium and LOS would be reduced by protocols such as orienting communication (ie, orientation and engaged conversation), oral and nutritional assistance (ie, brushing teeth, oral-facial exercise, and postoperative dietary education), and early mobilization.¹²⁻¹⁴ These protocols, initially developed in 2008, were modified from the Hospital Elder Life Program (HELP), which is cost-effective and has been disseminated widely.¹⁵⁻¹⁷ Our unique innovation was to select 3 core protocols and allow them to be delivered during daily care by trained nursing staff for feasibility and scalability.¹² For this cluster randomized clinical trial (RCT), we evaluated the effects of the modified Hospital Elder Life Program (mHELP) on delirium incidence and LOS in a sample of older patients (≥65 years of age) undergoing major elective abdominal surgery, primarily for resection of malignant tumors. As a subgroup analysis, effects were stratified by type of abdominal surgery.

Key Points

Question Does a modified Hospital Elder Life Program reduce incident delirium and hospital length of stay in patients undergoing abdominal surgery?


Findings In this cluster randomized clinical trial of 377 older patients undergoing elective abdominal surgery, postoperative delirium occurred in fewer patients in the intervention group than in the control group. Hospital length of stay was also significantly shorter in the intervention group.

Meaning The modified Hospital Elder Life Program strongly may benefit older patients undergoing abdominal surgery, with significant reduction of delirium incidence and hospital length of stay.

Methods

Although cluster randomization is less efficient than individual randomization because outcomes can be correlated between clusters (often reflected as the intraclass correlation [ICC]),¹⁸ this design minimizes contamination among participants in different groups by ensuring that all participants in one room belong to the same group. Physicians and hospital staff (surgeons, residents, and nurses) at the study site were aware of a pending nursing intervention study but were masked to study hypothesis, group allocation, and specific protocols of mHELP. Moreover, outcome assessors were masked to group assignment, and room assignments were rerandomized every 20 patients to minimize potential unmasking of the randomization scheme. The trial protocol can be found in Supplement 1. This cluster RCT was approved by the Research Ethics Review Committee at the National Taiwan University Hospital and registered at clinicaltrials.gov.

Figure 1. CONSORT Flow Study Diagram



LOS indicates length of stay; mHELP, modified Hospital Elder Life Program.

Patient Selection

Consecutive older patients (≥65 years of age) admitted to two 36-bed gastrointestinal wards of a 2000-bed urban medical center in Taipei, Taiwan, were screened for enrollment from August 1, 2009, through October 31, 2012. Patients were enrolled if they met 2 criteria: scheduled for elective abdominal surgery and expected LOS longer than 6 days. Participants were cluster randomized to groups with an allocation ratio of 1:1 based on a computer-generated list. Cluster randomization by room was necessary because most patient units in Taiwan are double- or triple-occupancy rooms, threatening cross-contamination if patients were individually randomized. This randomization approach was facilitated by both gastrointestinal wards having the same layout: 6 single-occupancy rooms (3 each randomly assigned to the control and mHELP groups), 9 double-occupancy rooms (4 randomly assigned to the control group and 5 to the mHELP group), and 4 triple-occupancy rooms (2 randomly assigned to each group) (Figure 1). Participants in the 38 rooms formed 318 clusters during the 3-year study period. Written informed consent was obtained for every participant in the study, and all study data were deidentified.

Intervention and Usual Care

The intervention (mHELP) was implemented by a trained mHELP nurse (registered nurse who had 2 years of medical-surgical experience and who was trained on site for 1 month before the intervention start)¹² who did not assess any outcomes. The intervention consisted of the daily hospital-based mHELP comprising 3 core nursing protocols: orienting communication, oral and nutritional assistance, and early mobilization (eAppendix 1 in Supplement 2).¹² In addition to usual perioperative care (eAppendix 2 in Supplement 2), participants received all 3 mHELP protocols postoperatively as soon as they arrived in the inpatient ward, immediately after interim intensive care stays, and until hospital discharge. All protocols were tracked daily with adherence rated on a Likert-type scale from 0 (no adherence) to 3 (full implementation and adherence).

Usual care consisted of standard hospital care provided by surgeons, residents, nurses, and physical therapists (as needed) in the general surgery wards. All participants were encouraged to ambulate and did so as tolerated. The mHELP nurses did not provide services to participants assigned to the control group. However, the same attending physicians provided care to participants in the mHELP and control groups.

Study Data

Two outcome assessors specially trained for delirium assessment collected outcome data from Monday through Saturday. Presence of delirium was assessed by the Confusion Assessment Method¹⁹ based on a brief daily cognitive screen and interview to rate 4 core delirium symptoms. Participants were considered to have delirium if they had the first (acute onset and fluctuating course) and second (inattention) core symptoms and the third (altered consciousness) or fourth (disorganized thinking) symptom. The Confusion Assessment Method is a widely used, standardized method for identifying delirium that has a sensitivity of 94% (95% CI, 91%-97%) and a specificity of 89% (95% CI, 85%-94%) compared with clinical expert ratings and an interrater reliability of 0.70 to 1.00.²⁰ Changes in mental status were also solicited from family members or nurses. The outcome assessors did not communicate with the mHELP nurses and were masked to participants' intervention status.

Patient characteristics obtained from in-person interviews included age, sex, and educational level. Baseline clinical factors included presurgical Charlson comorbidity index (higher scores indicate greater mortality risk),²¹ presurgical cognitive status measured using the Mini-Mental State Examination (score range, 0-30; 30 indicates no impairment),²² functional status measured using the Barthel Index (score range, 0-100; 100 indicates total independence),²³ nutritional status measured using the Mini-Nutritional Assessment (score range, 0-30; 30 indicates normal status),²⁴ and depressive status measured using the Geriatric Depression Scale Short Form (score range, 0-15; 15 indicates depression).²⁵ Other clinical data abstracted from medical records included diagnosis (gastric cancer, pancreatic or periampullary cancer, colorectal cancer, or other), malignant tumor (yes/no), tumor stage (0 to IV), type of surgery (total or subtotal gastrectomy; right hemicolec-

lectomy; left hemicolectomy, lower anterior resection, or anterior resection; pancreaticoduodenectomy; or other), duration of surgery (minutes), laparoscopic surgery (yes/no), intensive care unit (ICU) admission (yes/no), and length of ICU stay (days). The LOS data were abstracted from the medical record at discharge.

Statistical Analysis

Data were analyzed using an intention-to-treat approach. All analyses were performed with SAS statistical software, version 9.3 (SAS Institute Inc) and R software, version 3.2.1 (R Foundation for Statistical Computing). Sample characteristics were compared by treatment group at baseline. Data were reported as number (percentage), mean (SD), or median (interquartile range [IQR]) when not normally distributed.

An important feature of cluster RCTs is the extent of within-cluster correlation for end points. The ICC, defined as the ratio of between-cluster variance to total variance, refers to the proportion of variance attributed to the cluster level. The ICC and its 95% CI were calculated for each outcome using the ICCest function in the R software ICC, which adopted the variance components from a 1-way analysis of variance for the calculation.²⁶ Of note, all ICCs for each outcome (eAppendix 3 in Supplement 2) were not significantly different from 0 and some were even less than 0, suggesting that the true ICCs are small and adjustment for cluster effect is not indicated.²⁷ We thus analyzed treatment effects using standard statistical methods not accounting for within-cluster correlation. Kaplan-Meier analysis and the log-rank test were further used to compare the cumulative incidence of delirium, defined as the probability that delirium would develop during hospitalization, between study groups. All statistical tests were 2-tailed, and $P < .05$ was considered to indicate statistical significance. To correct for multiple comparisons, the significance of the intervention effect for each of the 5 surgical types was assessed at Bonferroni-corrected $P = .01$ (0.05/5).²⁸

Results

Of 577 eligible patients, 377 (65.3%) were enrolled and randomly assigned to the mHELP ($n = 197$; mean [SD] age, 74.3 [5.8] years; 111 [56.4%] male) or control ($n = 180$; mean [SD] age, 74.8 [6.0] years; 103 [57.2%] male) group (Figure 1 and Table 1).²⁹ The mHELP and control groups did not differ significantly in terms of any baseline characteristics, including presurgical cognitive status or other functional measures. The primary indication for surgery was malignant tumor (178 [90.4%] for the mHELP group vs 165 [91.7%] for the control group; $P = .64$).

Intervention Adherence

Participants and family caregivers reported positive perceptions of the mHELP protocols. The median start time of the intervention protocols was postoperative day 1 (IQR, 1-3 days), with 120 of 196 participants (61.2%) starting by postoperative day 1 and 173 participants (88.3%) receiving mHELP components no later than postoperative day 3. The reason for the de-

Table 1. Participants' Baseline Characteristics by Group^a

| Characteristic | mHELP (n = 197) | Control (n = 180) | P Value ^b |
|---|-----------------|-------------------|----------------------|
| Age, mean (SD), y | 74.3 (5.8) | 74.8 (6.0) | .38 |
| Male sex | 111 (56.4) | 103 (57.2) | .95 ^c |
| Educational level | | | |
| Illiterate | 25 (12.7) | 29 (16.1) | .54 ^c |
| Elementary or middle school | 90 (45.7) | 84 (46.7) | |
| High school and above | 82 (41.6) | 67 (37.2) | |
| Presurgical Charlson comorbidity index, mean (SD) | 1.6 (1.9) | 1.5 (1.7) | .83 |
| Presurgical Charlson comorbidity index | | | |
| 0 | 67 (34.0) | 59 (32.8) | .55 ^c |
| 1 | 49 (24.9) | 51 (28.3) | |
| ≥2 | 81 (41.1) | 70 (38.9) | |
| Presurgical scores, mean (SD) | | | |
| Cognitive MMSE ^d | 27.0 (3.8) | 26.8 (3.1) | .61 |
| Functional BI ^e | 97.1 (10.1) | 97.7 (6.2) | .50 |
| Nutritional MNA ^f | 24.7 (3.7) | 24.5 (3.9) | .70 |
| Depressive GDS ^g | 2.5 (2.6) | 2.7 (2.8) | .49 |
| Diagnosis | | | |
| Gastric cancer | 39 (19.8) | 41 (22.8) | .78 ^c |
| Pancreatic or periampullary cancer | 28 (14.2) | 24 (13.3) | |
| Colorectal cancer | 111 (56.4) | 102 (56.7) | |
| Other ^h | 19 (9.6) | 13 (7.2) | |
| Malignant tumor | 178 (90.4) | 165 (91.7) | .64 ^c |
| Tumor stage ⁱ | | | |
| 0 | 2 (1.1) | 6 (3.6) | .24 ^c |
| I | 45 (25.3) | 52 (31.5) | |
| II | 54 (30.3) | 35 (21.2) | |
| III | 49 (27.5) | 51 (30.9) | |
| IV | 28 (15.7) | 21 (12.7) | |
| Type of surgery ^j | | | |
| Total or subtotal gastrectomy | 43 (21.9) | 43 (24.0) | .52 ^c |
| Right hemicolectomy | 32 (16.3) | 32 (17.9) | |
| Left hemicolectomy, LAR, or AR | 67 (34.2) | 67 (37.4) | |
| Pancreaticoduodenectomy | 25 (12.8) | 21 (11.7) | |
| Other ^k | 29 (14.8) | 16 (8.9) | |
| Duration of surgery, median (IQR), min | 195 (105) | 213 (98) | .10 |
| Laparoscopy | 84 (42.6) | 93 (51.6) | .10 ^c |
| ICU admission after surgery | 100 (50.8) | 98 (54.4) | .47 ^c |
| Length of ICU stay, mean (SD), d | 2.8 (6.5) | 2.4 (3.8) | .58 |

Abbreviations: AR, anterior resection; BI, Barthel Index; GDS, 15-item Geriatric Depression Scale; ICU, intensive care unit; IQR, interquartile range; LAR, lower anterior resection; mHELP, modified Hospital Elder Life Program; MMSE, Mini-Mental State Examination; MNA, Mini-Nutritional Assessment.

^a Data are presented as number (percentage) of study participants unless otherwise indicated.

^b Significance was determined by Mann-Whitney test unless otherwise indicated.

^c Significance determined by χ^2 test.

^d Scores range from 0 to 30, with higher scores indicating better cognitive status.

^e Scores range from 0 to 100, with higher scores indicating better independence in activities of daily living.

^f Scores range from 0 to 30, with higher scores indicating better nutritional status.

^g Scores range from 0 to 15, with higher scores indicating more depressive symptoms.

^h Diagnoses included splenic tumor, mesothelioma, gastrointestinal stromal tumor, pseudomyxoma peritonei duodenum tumor, distal common bile duct tumor, pancreatic tumor, colon poly, and fistula.

ⁱ n = 178 for the mHELP group and 165 for the control group.

^j n = 196 for the mHELP group and 179 for the control group.

^k Open splenectomy, transverse colon partial resection, Hartmann procedure with adhesiolysis and bladder lithotripsy, abdominoperineal resection, or laparoscopic debulking surgery.

lay in the remaining 23 participants (11.7%) receiving mHELP components later than postoperative day 3 was that their ICU stay was prolonged beyond 3 days. Nevertheless, overall adherence to the protocols was good; 166 participants (84.3%) had mean scores of 2 or higher (range, 0-3), indicating moderately good adherence. Mean adherence scores for orienting communication and early mobilization were slightly higher than for oral and nutritional assistance (2.6 and 2.5 vs 2.3). In total, participants in the mHELP group received a median of 7 days (IQR, 6-10 days) of the mHELP protocols, and the mean (SD) time spent with each participant per session was 34.1 (16.0) minutes (median, 30 minutes; IQR, 25-40 minutes). No adverse events or unintended effects were reported as intervention related in the mHELP group.

Effects on Delirium

During hospitalization, 40 cases (10.6%) of incident delirium occurred in both groups. In the group that received mHELP, delirium developed in 13 cases (6.6%), whereas the control group had 27 cases (15.1%) (Table 2). These differences were statistically significant with a relative risk of 0.44 for delirium (95% CI, 0.23-0.83; $P = .008$), demonstrating a risk reduction of 56%. In absolute terms, the number of cases needed to treat to prevent 1 case of delirium was 11.8. The mHELP also had significant effects for cumulative incidence of delirium ($\chi^2 = 5.87, P = .02$) (Figure 2). Stratified by surgical type, participants who underwent total or subtotal gastrectomy and received mHELP had reduced delirium (1 [2.3%] in the mHELP group vs 8 [18.6%] in the control group; $P = .03$).

Table 2. Delirium and Length of Hospital Stay Outcomes

| Characteristic | mHELP | Control | P Value ^a |
|--|--------------|---------------|----------------------|
| Delirium, No./total No. (%) ^b | 13/196 (6.6) | 27/179 (15.1) | .008 ^c |
| Total or subtotal gastrectomy | 1/43 (2.3) | 8/43 (18.6) | .03 ^d |
| Right hemicolectomy | 1/32 (3.1) | 2/32 (6.3) | >.99 ^d |
| Left hemicolectomy, LAR, or AR | 6/67 (9.0) | 10/67 (14.9) | .43 ^d |
| Pancreaticoduodenectomy | 2/25 (8.0) | 6/21 (28.6) | .12 ^d |
| Other ^d | 3/29 (10.3) | 1/16 (6.3) | >.99 ^d |
| Length of stay, median (IQR), d ^e | 12.0 (6) | 14.0 (9) | .04 |
| Total or subtotal gastrectomy | 12.0 (5) | 18.0 (17) | <.001 |
| Right hemicolectomy | 12.0 (4) | 13.0 (5.5) | .12 |
| Left hemicolectomy, LAR, or AR | 12.0 (6) | 12.0 (5) | .79 |
| Pancreaticoduodenectomy | 16.0 (12) | 25.5 (25) | .28 |
| Other ^f | 12.0 (15) | 13.5 (7.5) | .95 |

Abbreviations: AR, anterior resection; IQR, interquartile range; LAR, lower anterior resection; mHELP, modified Hospital Elder Life Program.

^a Significance was determined by Mann-Whitney test unless indicated otherwise. Significance of the intervention effect for each of the 5 surgical types was assessed at the Bonferroni-corrected *P* value of .01 (0.05/5).

^b *n* = 196 for the mHELP group and 179 for the control group.

^c Significance determined by χ^2 test.

^d Significance determined by Fisher exact test.

^e *n* = 192 for the mHELP group and 176 for the control group.

^f Procedures such as open splenectomy, transverse colon partial resection, Hartmann procedure with adhesiolysis and bladder lithotripsy, abdominoperineal resection, and laparoscopic debulking surgery.

Effects on LOS


The mHELP and control groups differed significantly in median LOS (12.0 vs 14.0 days; *P* = .04) (Table 2). Stratified by surgical type, participants who underwent total or subtotal gastrectomy had significantly shorter LOS (12.0 vs 18.0 days; *P* < .001) with mHELP. Delayed implementation of mHELP components in 23 participants (11.7%) was attributable to a prolonged ICU stay of 3 days or longer. In this mHELP subgroup, delirium incidence was lower than that in the control subgroup (4 of 23 [17.4%] vs 6 of 20 [30.0%]), a difference that did not reach significance (*P* = .47). Moreover, LOS did not differ significantly between these subgroups (21.0 vs 21.0 days; *P* = .80).

Discussion

The mHELP strongly benefitted older patients undergoing abdominal surgery for resection of malignant tumor, with significant reduction of delirium incidence by 56% and hospital LOS by 2 days. As shown in Figure 2, development of delirium is not only delayed but also reduced for patients who received mHELP. Stratified by surgical type, patients who underwent gastrectomy benefited more from mHELP, with a 6-day shorter LOS than in the control group (12.0 vs 18.0 days; *P* < .001). This subgroup also experienced a trend toward reduced delirium incidence. The mechanism for this greater benefit in patients undergoing gastrectomy is unclear, requiring further research to understand factors that may magnify or attenuate the mHELP effects and to define the effect of mHELP in various surgical procedures.

Consistent with our RCT findings, a 14-study meta-analysis¹⁷ found that multicomponent, nonpharmacologic interventions including at least 2 to 6 components (ie, cognition, mobilization, hydration, hearing, vision, and sleep-wake cycle) in 4 randomized or matched trials (mostly medical inpa-

Figure 2. Cumulative Incidence of Delirium by Group



The cumulative incidence of delirium was defined as the probability of the development of delirium during hospitalization. Data on patients were censored at the time of discharge or death. The difference between the groups was significant ($\chi^2 = 5.87$; *P* = .02 by the log-rank test). Because of the smaller sample sizes, the figure does not extend beyond 18 days. mHELP indicates modified Hospital Elder Life Program.

tients; one focusing on surgical patients) effectively reduced incident delirium by 44% with a trend toward reducing LOS. The mHELP targets similar components (orienting communication, oral and nutritional assistance, and early mobilization) with a unique extension to brushing teeth and oral-facial exercise to improve dry mouth and swallowing efficacy, thus facilitating oral intake. We postulated that increasing older patients' attention to and engagement with the postoperative recovery environment,³⁰ increasing their swallowing efficacy and nutritional and fluid repletion,^{23,26,31} and augmenting physical activity^{32,33} would prevent delirium and reduce LOS. Future research is needed to elucidate the mechanisms of the intervention effect; possible research areas include neu-

ropsychological measures, such as executive functioning and attention; physiology of swallowing efficacy; nutritional and fluid parameters; or inflammatory markers.

We note that other delirium prevention approaches for older hospitalized adults have included proactive geriatric consultation,³⁴ training family members,^{35,36} sustained education,³⁷⁻³⁹ and single-component interventions, such as bright light, music therapy, and use of software to detect medications that may cause delirium.⁴⁰ Not all studies^{35,36,38,39,41} included surgical patients or documented efficacy in reducing delirium incidence. An important issue noted by researchers in most of these studies^{35,39,41} was that assuring adherence to the interventions was a key factor for success and was not always achievable across settings.⁴¹

Indeed, the 3 mHELP protocols might seem commonsensical, yet the key to their effectiveness may lie in their consistent daily application.⁴² In this study, we had a full-time-equivalent trained mHELP nurse to consistently deliver all 3 protocols to 196 patients, spending approximately 30 minutes with each patient daily. Thus, with an additional 30 minutes of nursing time per older patient, mHELP reduced delirium by 56% and shortened LOS by 2 days, which will greatly reduce medical costs. By extrapolation, older patients in the United States had 7.96 million surgical hospital stays in 2012, with a mean cost of \$11 600 per stay.⁴³ Thus, mHELP could have prevented approximately 674 576 cases of delirium in the surgical service in 2012, resulting in a Medicare cost savings of approximately \$10 000 per case⁴⁴ or \$6.7 billion for the year.^{4,5} By cutting 2 days from LOS (of 14 days in controls; a 14% reduction), implementation of mHELP could have saved \$1624 per hospital stay or \$12.9 billion per year in Medicare costs for the hospital stay.

Limitations

Several caveats about this study are worthy of comment. First, we did not adjust for the cluster effect because of very small ICCs, indicating weak between-cluster correlations for each outcome. To gain efficiency, future trials may use individual randomization instead of cluster randomization. Second, with a sample size of 377, post hoc analysis indicated that our study was powered at 81% for delirium and 80% for LOS to detect group differences as a whole but was underpowered for subgroup analyses by surgical type. Third, 9 of 377 participants (2.4% attrition rate, including 3 deaths and 6 dropouts) had

missing outcome values, which might have biased the study findings. However, this bias was likely minimized by attrition rates not differing significantly between the intervention and control groups (2.5% vs 2.2%). Fourth, participants from both groups received care from the same surgeons and nurses; that is, some participants in the control group may have received mHELP components through crossover (contamination) effects. However, the effect of this contamination would have underestimated the mHELP effects. Fifth, we did not collect data on postoperative complications, which are important risk factors for delirium and might have also been affected by mHELP and contributed to the study findings. Sixth, our trial was conducted without an enhanced recovery after surgery (ERAS) program that involved epidural or regional anesthesia, minimally invasive techniques, fluid and pain management, and aggressive postoperative rehabilitation.⁴⁵ Although this omission may limit generalizability to centers using ERAS, mHELP may still present important advantages. For medical centers unable to initiate a full ERAS program, mHELP may be considered to be a useful starting point to advance care for vulnerable older patients. Moreover, for centers with ERAS already implemented, mHELP provides feasible, structured, postoperative care protocols that target cognition, nutrition, and ambulation to augment the ERAS program and enhance recovery.

Conclusions

Delirium, which is recognized as the most common surgical complication in older patients, has been associated with increased morbidity and mortality, prolonged hospital stays, higher medical costs, and greater likelihood of institutionalization.^{6,46} Older patients undergoing major abdominal surgery for resection of malignant tumor had markedly reduced rates of incident delirium and shorter LOS when they received mHELP, which included 3 nurse-administered protocols: orienting communication, oral and nutritional assistance, and early mobilization. The key to the effectiveness of the 3 mHELP components is their consistent and daily application, with high adherence rates.⁴² Medical centers that want to advance postoperative care for older patients might consider mHELP as a highly effective starting point for delirium prevention.

ARTICLE INFORMATION

Accepted for Publication: March 9, 2017.

Published Online: May 24, 2017.

doi:10.1001/jamasurg.2017.1083

Author Affiliations: Department of Nursing, College of Medicine, National Taiwan University, Taipei, Taiwan (C.-H. Chen, Li); Department of Nursing, National Taiwan University Hospital, Taipei, Taiwan (C.-H. Chen); Department of Nursing, Sijihh Cathy General Hospital, New Taipei City, Taiwan (Li); Department of Surgery, College of Medicine, National Taiwan University, Taipei, Taiwan (Liang, Lai, B.-R. Lin, J. Huang, C.-Y. Yang, Tien, C.-N. Chen, M.-T. Lin); Department of Surgery,

National Taiwan University Hospital, Taipei, Taiwan (Liang, Lai, B.-R. Lin, J. Huang, C.-Y. Yang, Tien, C.-N. Chen, M.-T. Lin); Institute of Statistics, National Chiao Tung University, Hsinchu, Taiwan (Purnomo, G.-H. Huang); Taiwan Center for Disease Control, Taipei, Taiwan (Y.-T. Yang); Department of Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts (Inouye); Institute for Aging Research, Hebrew Senior-Life, Boston, Massachusetts (Inouye).

Author Contributions: Dr C.C.-H. Chen had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: C.C.-H. Chen, Liang, Lai, B.-R. Lin, J. Huang, C.-N. Chen, M.-T. Lin, Inouye. **Acquisition, analysis, or interpretation of data:** C.C.-H. Chen, Li, Liang, Purnomo, Y.-T. Yang, C.-Y. Yang, Tien, M.-T. Lin, G.-H. Huang, Inouye. **Drafting of the manuscript:** C.C.-H. Chen, Liang, Purnomo, Y.-T. Yang, C.-N. Chen, M.-T. Lin, G.-H. Huang.

Critical revision of the manuscript for important intellectual content: C.C.-H. Chen, Li, Liang, Lai, B.-R. Lin, J. Huang, C.-Y. Yang, Tien, M.-T. Lin, G.-H. Huang, Inouye.

Statistical analysis: C.C.-H. Chen, Li, Liang, Purnomo, Y.-T. Yang, M.-T. Lin, G. Huang.

Obtained funding: C.C.-H. Chen, Liang, M.-T. Lin.

Administrative, technical, or material support: C.C.-H. Chen, Li, Liang, Lai, B.-R. Lin, J. Huang, C.-Y. Yang, Tien, C.-N. Chen, M.-T. Lin, Inouye.
Study supervision: C.C.-H. Chen, Liang, Tien, M.-T. Lin, Inouye.

Conflict of Interest Disclosures: None reported.

Funding/Support: This study was supported in part by grants 98-2314-B-002-113-MY3 from the Ministry of Science and Technology and NHRI-EX-9820PC from the National Health Research Institute in Taiwan (Dr C.C.-H. Chen). Dr Inouye's time was covered in part by grants R24AG054259, P01AG031720, K07AG041835, and R01AG044518 from the National Institute on Aging. Dr Inouye holds the Milton and Shirley F. Levy Family Chair.

Role of the Funder/Sponsor: The funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

REFERENCES

- Rudolph JL, Marcantonio ER. Postoperative delirium: acute change with long-term implications. *Anesth Analg*. 2011;112(5):1202-1211.
- van Bree SH, Vlug MS, Bemelman WA, et al. Faster recovery of gastrointestinal transit after laparoscopy and fast-track care in patients undergoing colonic surgery. *Gastroenterology*. 2011;141(3):872-880.e1, 4.
- Augestad KM, Delaney CP. Postoperative ileus: impact of pharmacological treatment, laparoscopic surgery and enhanced recovery pathways. *World J Gastroenterol*. 2010;16(17):2067-2074.
- Inouye SK, Westendorp RGJ, Saczynski JS. Delirium in elderly people. *Lancet*. 2014;383(9920):911-922.
- Leslie DL, Inouye SK. The importance of delirium: economic and societal costs. *J Am Geriatr Soc*. 2011;59(suppl 2):S241-S243.
- Gleason LJ, Schmitt EM, Kosar CM, et al. Effect of delirium and other major complications on outcomes after elective surgery in older adults. *JAMA Surg*. 2015;150(12):1134-1140.
- Cavallari M, Hshieh TT, Guttman CR, et al; SAGES Study Group. Brain atrophy and white-matter hyperintensities are not significantly associated with incidence and severity of postoperative delirium in older persons without dementia. *Neurobiol Aging*. 2015;36(6):2122-2129.
- Silverstein JH. Cognition, anesthesia, and surgery. *Int Anesthesiol Clin*. 2014;52(4):42-57.
- Young J, Inouye SK. Delirium in older people. *BMJ*. 2007;334(7598):842-846.
- Scholz AF, Oldroyd C, McCarthy K, Quinn TJ, Hewitt J. Systematic review and meta-analysis of risk factors for postoperative delirium among older patients undergoing gastrointestinal surgery. *Br J Surg*. 2016;103(2):e21-e28.
- Siddiqi N, House AO, Holmes JD. Occurrence and outcome of delirium in medical in-patients: a systematic literature review. *Age Ageing*. 2006;35(4):350-364.
- Chen CC, Saczynski J, Inouye SK. The modified Hospital Elder Life Program: adapting a complex intervention for feasibility and scalability in a surgical setting. *J Gerontol Nurs*. 2014;40(5):16-22.
- Chen CC, Lin MT, Tien YW, Yen CJ, Huang GH, Inouye SK. Modified Hospital Elder Life Program: effects on abdominal surgery patients. *J Am Coll Surg*. 2011;213(2):245-252.
- Chen CC, Chen CN, Lai IR, Huang GH, Saczynski JS, Inouye SK. Effects of a modified Hospital Elder Life Program on frailty in individuals undergoing major elective abdominal surgery. *J Am Geriatr Soc*. 2014;62(2):261-268.
- Inouye SK, Baker DI, Fugal P, Bradley EH; HELP Dissemination Project. Dissemination of the Hospital Elder Life Program: implementation, adaptation, and successes. *J Am Geriatr Soc*. 2006;54(10):1492-1499.
- Rubin FH, Neal K, Fenlon K, Hassan S, Inouye SK. Sustainability and scalability of the Hospital Elder Life Program at a community hospital. *J Am Geriatr Soc*. 2011;59(2):359-365.
- Hshieh TT, Yue J, Oh E, et al. Effectiveness of multicomponent nonpharmacological delirium interventions: a meta-analysis. *JAMA Intern Med*. 2015;175(4):512-520.
- Campbell MJ. Challenges of cluster randomized trials. *J Comp Eff Res*. 2014;3(3):271-281.
- Inouye SK, van Dyck CH, Alessi CA, Balkin S, Siegel AP, Horwitz RI. Clarifying confusion: the confusion assessment method: a new method for detection of delirium. *Ann Intern Med*. 1990;113(12):941-948.
- Wei LA, Fearing MA, Sternberg EJ, Inouye SK. The Confusion Assessment Method: a systematic review of current usage. *J Am Geriatr Soc*. 2008;56(5):823-830.
- Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373-383.
- Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975;12(3):189-198.
- Rofes L, Arreola V, Romea M, et al. Pathophysiology of oropharyngeal dysphagia in the frail elderly. *Neurogastroenterol Motil*. 2010;22(8):851-858, e230.
- Guigoz Y, Vellas B, Garry PJ. Assessing the nutritional status of the elderly: the Mini Nutritional Assessment as part of the geriatric evaluation. *Nutr Rev*. 1996;54(1, pt 2)(suppl 2):S59-S65.
- D'Ath P, Katona P, Mullan E, Evans S, Katona C. Screening, detection and management of depression in elderly primary care attenders, I: the acceptability and performance of the 15 item Geriatric Depression Scale (GDS15) and the development of short versions. *Fam Pract*. 1994;11(3):260-266.
- Marcantonio ER. Postoperative delirium: a 76-year-old woman with delirium following surgery. *JAMA*. 2012;308(1):73-81.
- McDonald JH. *Handbook of Biological Statistics*. 3rd ed. Baltimore, MD: Sparky House Publishing; 2014.
- Xu Y, Lee CF, Cheung YB. Analyzing binary outcome data with small clusters: a simulation study. *Commun Stat Simul Comput*. 2014;43(7):1771-1782.
- Moher D, Hopewell S, Schulz KF, et al. CONSORT 2010 explanation and elaboration: updated guidelines for reporting parallel group randomised trials. *BMJ*. 2010;340:c869.
- Cheng CM, Chiu MJ, Wang JH, et al. Cognitive stimulation during hospitalization improves global cognition of older Taiwanese undergoing elective total knee and hip replacement surgery. *J Adv Nurs*. 2012;68(6):1322-1329.
- Lim P, Morris OJ, Nolan G, Moore S, Draganic B, Smith SR. Sham feeding with chewing gum after elective colorectal resectional surgery: a randomized clinical trial. *Ann Surg*. 2013;257(6):1016-1024.
- Ertek S, Cicero A. Impact of physical activity on inflammation: effects on cardiovascular disease risk and other inflammatory conditions. *Arch Med Sci*. 2012;8(5):794-804.
- Dainese R, Serra J, Azpiroz F, Malagelada JR. Influence of body posture on intestinal transit of gas. *Gut*. 2003;52(7):971-974.
- Marcantonio ER, Flacker JM, Wright RJ, Resnick NM. Reducing delirium after hip fracture: a randomized trial. *J Am Geriatr Soc*. 2001;49(5):516-522.
- Rosenbloom-Brunton DA, Henneman EA, Inouye SK. Feasibility of family participation in a delirium prevention program for hospitalized older adults. *J Gerontol Nurs*. 2010;36(9):22-33.
- Martinez FT, Tobar C, Beddings CI, Vallejo G, Fuentes P. Preventing delirium in an acute hospital using a non-pharmacological intervention. *Age Ageing*. 2012;41(5):629-634.
- Lundström M, Olofsson B, Stenvall M, et al. Postoperative delirium in old patients with femoral neck fracture: a randomized intervention study. *Aging Clin Exp Res*. 2007;19(3):178-186.
- Tabet N, Hudson S, Sweeney V, et al. An educational intervention can prevent delirium on acute medical wards. *Age Ageing*. 2005;34(2):152-156.
- Milisen K, Foreman MD, Abraham IL, et al. A nurse-led interdisciplinary intervention program for delirium in elderly hip-fracture patients. *J Am Geriatr Soc*. 2001;49(5):523-532.
- Abraha I, Trotta F, Rimland JM, et al. Efficacy of non-pharmacological interventions to prevent and treat delirium in older patients: a systematic overview. The SENATOR project ONTOP Series. *PLoS One*. 2015;10(6):e0123090.
- Young J, Cheater F, Collinson M, et al. Prevention of delirium (POD) for older people in hospital: study protocol for a randomised controlled feasibility trial. *Trials*. 2015;16:340.
- Greysen SR. Delirium and the "know-do" gap in acute care for elders. *JAMA Intern Med*. 2015;175(4):521-522.
- Weiss A, Elixhauser A. Overview of hospital stays in the United States, 2012. <http://www.hcup-us.ahrq.gov/reports/statbriefs/sb180-Hospitalizations-United-States-2012.pdf>. Accessed June 29, 2016.
- Leslie DL, Zhang Y, Bogardus ST, Holford TR, Leo-Summers LS, Inouye SK. Consequences of preventing delirium in hospitalized older adults on nursing home costs. *J Am Geriatr Soc*. 2005;53(3):405-409.

45. Nicholson A, Lowe MC, Parker J, Lewis SR, Alderson P, Smith AF. Systematic review and meta-analysis of enhanced recovery programmes in surgical patients. *Br J Surg*. 2014;101(3):172-188.

46. American Geriatrics Society Expert Panel on Postoperative Delirium in Older Adults. Postoperative delirium in older adults: best practice

statement from the American Geriatrics Society. *J Am Coll Surg*. 2015;220(2):136-48.e1.

 Invited Commentary

Interventions to Reduce Postoperative Delirium

Aligning Surgical Care With Patients' Needs and Priorities

Pasithorn A. Suwanabol, MD; Daniel B. Hinshaw, MD

In this issue of *JAMA Surgery*, Chen and colleagues¹ report a cluster randomized clinical trial of a modified Hospital Elder Life Program encompassing multicomponent nonpharmacologic interventions to reduce postoperative delirium in older adults undergoing major abdominal surgery. The interventions include orienting communications, oral and nutritional assistance, and early mobilization in addition to usual postoperative care, together requiring 30 additional minutes of care per day yet significantly reducing the incidence of postoperative delirium and length of hospital stay. The authors report that these simple interventions with minimal risk can have profound effects that are scalable and may be easily incorporated into existing postoperative protocols.

Delirium affects 13% to 50% of surgical patients and is estimated to cost \$152 billion per year in the United States.^{2,3} Postoperative delirium is associated with increased length of hospital stay, rates of nonhome discharge, and mortality rates.³ Furthermore, delirium can affect postoperative function and long-term prognosis in addition to leading to a substantially higher risk of persistent cognitive decline and the development of dementia and depression.⁴ Cognitive impairments affect the ability to care for oneself, restrict social functioning, and decrease decision-making capacity. Most recently, Pusswald et al⁵ reported an association between self-reported impairments in cognition and reduced health-related quality of life. A significant focus has been made on

modifiable factors to improve postoperative outcomes in older adults, yet few studies, at least in the general surgery literature, examine postoperative interventions to reduce the incidence of delirium in this patient population.⁶

This study highlights not only a feasible and effective intervention but also notably outcome measures that are most important to patients. Delirium, subsequent cognitive decline, and potential for dementia are distressing to patients and families, leading to decreased or loss of functional ability (including threatened loss of independent living), depressive symptoms, and poorer quality of life. It is critical that we continue to examine these long-term outcomes of surgery on older adults and find measures to reduce these burdensome effects. Hospital Elder Life Programs have been implemented at more than 200 sites in the United States and worldwide, with an overall reduction in postoperative delirium by 30%.⁷ However, this study and most available delirium intervention literature fail to address the surgical intensive care unit patient population. Future studies in these settings are warranted, which may further mitigate the incidence of delirium and its sequelae. Nonetheless, interventions such as these not only reduce health care cost but also improve patient quality of life and address patient priorities that may not be measured by typical surgical quality metrics, such as death and complications. The surgical community should take notice of this important work because it may serve as a cost-effective model for achieving outcomes that are meaningful to surgeons and their patients.

ARTICLE INFORMATION

Author Affiliations: Department of Surgery, University of Michigan, Ann Arbor (Suwanabol, Hinshaw); Palliative Care Program, University of Michigan Geriatrics Center, Ann Arbor (Hinshaw).

Corresponding Author: Pasithorn A. Suwanabol, MD, Department of Surgery, University of Michigan, 1500 E Medical Center Dr, Taubman Center, Ste 2124, Ann Arbor, MI 48019 (pasuwan@med.umich.edu).

Published Online: May 24, 2017.
doi:10.1001/jamasurg.2017.1084

Conflict of Interest Disclosures: None reported.

REFERENCES

1. Chen CC-H, Li H-C, Liang J-T, et al. Effect of a modified Hospital Elder Life Program on delirium

and length of hospital stay in patients undergoing abdominal surgery: a cluster randomized clinical trial [published online May 24, 2017]. *JAMA Surg*. doi:10.1001/jamasurg.2017.1083

2. Inouye SK, Westendorp RG, Saczynski JS. Delirium in elderly people. *Lancet*. 2014;383(9920):911-922.

3. Leslie DL, Marcantonio ER, Zhang Y, Leo-Summers L, Inouye SK. One-year health care costs associated with delirium in the elderly population. *Arch Intern Med*. 2008;168(1):27-32.

4. Inouye SK, Marcantonio ER, Kosar CM, et al. The short-term and long-term relationship between delirium and cognitive trajectory in older surgical patients. *Alzheimers Dement*. 2016;12(7):766-775.

5. Pusswald G, Tropper E, Kryspin-Exner I, et al. Health-related quality of life in patients with subjective cognitive decline and mild cognitive impairment and its relation to activities of daily living. *J Alzheimers Dis*. 2015;47(2):479-486.

6. Siddiqi N, Harrison JK, Clegg A, et al. Interventions for preventing delirium in hospitalised non-ICU patients. *Cochrane Database Syst Rev*. 2016;3:CD005563.

7. Zauber TS, Murphy K, Rizzuto L, et al. Quality improvement and cost savings with multicomponent delirium interventions: replication of the Hospital Elder Life Program in a community hospital. *Psychosomatics*. 2013;54(3):219-226.